

Infographic. Clinical recommendations for return to play during the COVID-19 pandemic

Herbert Löllgen,^{1,2} Norbert Bachl,^{1,3,4,5} Theodora Papadopoulou,^{1,3,6,7} Andrew Shafik,⁷ Graham Holloway,⁷ Karin Vonbank,⁸ Nigel Edward Jones,^{7,9} Xavier Bigard,^{1,10} David Niederseer ,¹¹ Joachim Meyer,^{12,13} Borja Muniz-Pardos,¹⁴ Andre Debruyne,^{1,3} Petra Zupet,^{1,15} Jürgen M Steinacker ,^{1,3,16} Bernd Wolfarth,^{3,17} James Lee John Bilzon ,^{3,7,18} Anca Ionescu,^{1,19} Michiko Dohi ,^{3,20} Jeroen Swart,^{3,21} Victoriya Badtieva,^{3,22,23} Irina Zelenkova,^{14,22} Maurizio Casasco,^{1,3,24} Michael Geistlinger,^{3,25} Luigi Di Luigi,^{3,26} Nick Webbhorn,²⁷ Patrick Singleton,²⁸ Mike Miller,²⁸ Fabio Pigozzi,^{1,3,29,30} Yannis P Pitsiladis ^{1,3,31}

COVID-19 AND RETURN TO PLAY

The world of sport has recently returned to training and competition following suspension due to the COVID-19 pandemic. It is concerning that a number of athletes have tested positive for COVID-19 after returning to competition.¹ Numerous authors have attempted to address return to play given the importance and complexity of the issue, with notable attention on possible cardiac implications.²⁻⁶

SCOPE OF THE INFOGRAPHIC

The specific recommendations shown in the present infographic (figure 1) have been generated by a panel of international experts and represent a compilation of the numerous approaches used to inform resumption of regular sports during the COVID-19 pandemic. Despite the different regulations around the world and the particular characteristics of each sport, it is essential to provide informative, consistent and specific guidance for safe return to training and competition at this most difficult time.

RECOMMENDATIONS

COVID-19 is a systemic disease affecting the endothelium and all organs, especially the heart and lungs with more severity and clinical significance. Therefore, our recommendations for safe return to sport are based on a comprehensive clinical evaluation (ie, cardiac and respiratory) and have been designed for six different groups according to the level of exposure to COVID-19. These groups are based on a previous report by Phelan *et al.*,³ which are applicable to both leisure time and professional athletes. It is imperative that group allocation is determined beforehand through a medical examination performed in cooperation with a respiratory physician and/or cardiologist. Specific recommendations for each group are provided based on current evidence and expert opinion (see table 1).

SPECIAL CONSIDERATIONS

The resumption of sporting activity should be determined on a case-by-case basis and consider the individual situation of the athlete, including pre-existing conditions, as well as the type of sport and the risk of infection from other

athletes (eg, increased risk in contact/team sports). Ideally, the final decision to return to play will be based on the results of the individual assessment in consultation with the sport and exercise medicine physician, specialists in pulmonary medicine and sport cardiologists (or




Figure 1 Recommendations for return to sport during the COVID-19 pandemic. BNP, brain natriuretic peptide; CRP, C-reactive protein.

Table 1 Clinical recommendations for testing individual groups

Recommendations	Groups					
	1	2	3	4	5	6
Tested randomly or by finding antibodies. Only possible PPE or standard exam	X					
Need for usual check-up including ECG and echo		X				
Tested following contact with somebody who was positive or tested because having typical signs and symptoms and out of hospital treatment		X				
In-hospital treatment but quarantine at home			X	X		
For diagnosis only			X			
Treatment with close monitoring due to symptoms but normal ward with isolation and quarantine					X	
Need thorough preparticipation exam, CPET, echo and other necessary assessments			X	X		
Serious illness with difficult progression and intensive care unit without artificial ventilation					X	X
Artificial ventilation						X
Need thorough examination before discharge (eg, MRI)					X	X

CPET, cardiopulmonary exercise testing; PPE, pre-participation evaluation.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Löllgen H, Bachl N, Papadopoulou T, et al. *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2020-102985

Accepted 22 August 2020

Br J Sports Med 2020;0:1–2.
doi:10.1136/bjsports-2020-102985

ORCID iDs

- David Niederseer <http://orcid.org/0000-0003-3089-1222>
- Jürgen M Steinacker <http://orcid.org/0000-0001-6210-2449>
- James Lee John Bilzon <http://orcid.org/0000-0002-6701-7603>
- Michiko Dohi <http://orcid.org/0000-0002-1126-7849>
- Yannis P Pitsiladis <http://orcid.org/0000-0001-6210-2449>

REFERENCES

- 1 Fuller R. Novak Djokovic must 'feel responsibility' at coronavirus at Adria Tour, says Evans - BBC Sport. Available: <https://www.bbc.com/sport/tennis/53131665> [Accessed 27 Jun 2020].
- 2 Elliott N, Martin R, Heron N, et al. Infographic. Graduated return to play guidance following COVID-19 infection. *Br J Sports Med* 2020. doi:10.1136/bjsports-2020-102637. [Epub ahead of print: 22 Jun 2020].
- 3 Phelan D, Kim JH, Chung EH. A game plan for the resumption of sport and exercise after coronavirus disease 2019 (COVID-19) infection. *JAMA Cardiol* 2020. doi:10.1001/jamacardio.2020.2136. [Epub ahead of print: 13 May 2020].
- 4 EFSMA recommendations for sports and exercise medicine physicians: resumption of regular sports activity after COVID-19 pandemic. Available: <https://www.efsm.eu/newsevents/news-events/247-efsm-recommendations-for-sports-and-exercise-medicine-physicians-resumption-of-regular-sports-activity-after-covid-19-pandemic.html> [Accessed 27 Jun 2020].
- 5 Baggish A, Drezner JA, Kim J, et al. Resurgence of sport in the wake of COVID-19: cardiac considerations in competitive athletes. *Br J Sports Med* 2020. doi:10.1136/bjsports-2020-102516. [Epub ahead of print: 19 Jun 2020].
- 6 Does H, Cardin N. Return to play after COVID-19: a sport cardiologist's view. *Br J Sports Med*;395:bjsports-2020-102482.
- 7 Carmody S, Murray A, Borodina M, et al. When can professional sport recommence safely during the COVID-19 pandemic? Risk assessment and factors to consider. *Br J Sports Med* 2020;54:946–8.
- 8 Niess AM, Bloch W, Friedmann-Bette B, et al. Position stand: return to sport in the current coronavirus pandemic. *Dtsch Z SportMed*2020;71:E1–2.

extended multidisciplinary team), coaches and training specialists. After a prolonged period of interruption in sport caused by more severe health issues, increments in training should be gradual, individualised and supervised by periodic monitoring of cardiac and respiratory markers to detect potential signs and symptoms. Although the present recommendations are focused primarily on pulmonary and cardiac assessments, given the wider health-related issues caused by COVID-19, more comprehensive evaluation including psychological status, examination of muscular function and other suspected collateral issues should be considered to provide more detailed and thorough information for safe return to play.

FURTHER ESSENTIAL READING

For more detailed description of the risk factors and symptoms that could inform the decision to return to sport as safely as possible, consult Baggish et al,⁵ Carmody et al⁷ and Niess et al.⁸

- ¹European Federation of Sports Medicine Associations (EFSMA), Lausanne, Switzerland
- ²Cardiology Practice, Remscheid, Germany
- ³International Federation of Sports Medicine (FIMS), Lausanne, Switzerland
- ⁴Institute of Sports Science, University of Vienna, Vienna, Austria
- ⁵Austrian Institute of Sports Medicine, Vienna, Austria
- ⁶Lower Limbs-ADMR Hip & Groin, Defence Medical Rehabilitation Centre Headley Court, Loughborough, UK
- ⁷British Association of Sport and Exercise Medicine, Doncaster, UK
- ⁸Department of Pulmology, Medical University of Vienna, Vienna, Austria
- ⁹University of Liverpool, Liverpool, UK
- ¹⁰Union Cycliste Internationale (UCI), Aigle, Switzerland
- ¹¹Department of Cardiology, University Heart Center Zurich, Zurich, Switzerland
- ¹²German Respiratory Society (DGP), Berlin, Germany
- ¹³Lungenzentrum (Bogenhausen-Harlaching), München Klinik, Munich, Germany

- ¹⁴GENUD (Growth, Exercise, Nutrition and Development), Department of Physiatry and Nursing, University of Zaragoza, Zaragoza, Spain
- ¹⁵Institute for Medicine and Sport, Ljubljana, Slovenia
- ¹⁶Division of Sports and Rehabilitation Medicine, Ulm University Hospital, Ulm, Germany
- ¹⁷Department of Sport Medicine, Humboldt University and Charité University School of Medicine, Berlin, Germany
- ¹⁸Department for Health, University of Bath, Bath, UK
- ¹⁹Sports Medicine, "Carol Davila" University of Medicine, Bucharest, Romania
- ²⁰Sport Medical Center, Japan Institute of Sports Sciences, Tokyo, Japan
- ²¹UCT Research Unit for Exercise Science and Sports Medicine, University of Cape Town, Cape Town, South Africa
- ²²I.M. Sechenov First Moscow State Medical University (Sechenov University), Ministry of Health of Russia, Moscow, Russian Federation
- ²³Moscow Research and Practical Centre for Medical Rehabilitation, Restorative and Sports Medicine, Moscow Healthcare Department, Moscow, Russian Federation
- ²⁴Italian Federation of Sports Medicine (FMSI), Rome, Italy
- ²⁵Department of Public, International and European Law, Unit of International Law, University of Salzburg, Salzburg, Austria
- ²⁶Movement, Human and Health Sciences, University of Rome Foro Italico, Rome, Italy
- ²⁷School of Sport and Service Management, University of Brighton, Eastbourne, UK
- ²⁸World Olympians Association, Lausanne, Switzerland
- ²⁹Sports Medicine Unit, University of Rome Foro Italico, Rome, Italy
- ³⁰FIFA Medical Center of Excellence, Villa Stuart Sport Clinic, Rome, Italy
- ³¹Collaborating Centre of Sports Medicine, University of Brighton, Eastbourne, UK

Correspondence to Professor Yannis P Pitsiladis, Collaborating Centre of Sports Medicine, University of Brighton, Eastbourne BN20 7SN, UK; Y.Pitsiladis@Brighton.ac.uk

Twitter Theodora Papadopoulou @Dora_Sportmed, James Lee John Bilzon @JBilzon, Michiko Dohi @rinsho and Patrick Singleton @PSingleton33

Contributors All authors contributed significantly and in line with the instructions to authors to merit publication.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.